

2016 Gaps in Care Adult Measures
QIP/CTC/BCBSRI/UHC/MSSP

	Condition	Payer	Measure	Description	Target
1	Diabetes	CTC UHC Comm	HbA1c Testing	Patients 18-75 with diabetes who had an HbA1c test during the measurement period.	94%
2	Diabetes	CTC QIP (P4P)	HbA1c Poor Control	The percentage of patients 18-75 years of age with diabetes whose most recent HbA1c level during the measurement period is >9.0% (poor control) or there is no HbA1c in the record.	
3	Diabetes	BCBS CTC MSSP QIP (P4P) UHC Comm UHC MA	HbA1c Control	The percentage of patients 18-75 years of age with diabetes with the most recent HbA1c <8.0% or ≤9.0% (control) during the measurement period.	72%
4	Diabetes	BCBS UHC Comm UHC MA	Diabetes Care - Kidney Disease Monitoring (Nephropathy Screening)	The percentage of members 18–75 years of age with diabetes who had medical attention for nephropathy or had a urine micro albumin test during the measurement period.	86%
5	Diabetes	BCBS MSSP UHC Comm UHC MA	Diabetic Care Eye Exam	The percentage of members 18–75 years of age with diabetes who had an eye exam (retinal or dilated) performed during the measurement period or a negative retinal exam in the prior measurement period.	74%
6	Diabetes	CTC QIP (P4P)	Blood Pressure Control	The percentage of patients 18-75 with diabetes whose most recent blood pressure reading value is less than 140/90.	80%
7	Prevention	CTC MSSP QIP (P4P)	Tobacco Use	Patients age 18 years and older seen by a physician a minimum of once per measurement period who were queried about tobacco use a minimum of once per measurement period.	95%
8	Prevention	CTC QIP (P4P)	Tobacco Cessation Intervention	Patients age 18 years and older seen by a physician a minimum of once per measurement period who have been identified as smokers or tobacco users who received cessation intervention – (advice to quit, counseling, referral to counseling, and/or pharmacologic therapy during the MP ¹).	85%
9	Prevention	BCBS CTC MSSP QIP (P4P) UHC Comm UHC MA	Adult Body Mass Index (BMI)	The percentage of patients 18–74 years of age who had an outpatient visit and who had their “latest” body mass index (BMI) documented during the measurement period and whose BMI is either in the normal range OR is above or below the normal range and have a documented follow up plan within the measurement period.	87%
10	Prevention	CTC QIP (P4P) MSSP UHC MA	Screening for Fall Risk	Percentage of patients aged 65 years and older who were screened for future fall risk at least once within the MP ¹ .	70%

¹ MP = Measurement Period

2016 Gaps in Care Adult Measures
QIP/CTC/BCBSRI/UHC/MSSP

	Condition	Payer	Measure	Description	Target
11	Hypertension	BCBS CTC MSSP QIP (P4P)	Blood Pressure Control	The percentage of patients 18-85 years of age who had a diagnosis of hypertension (HTN) and their BP was adequately controlled during the measurement period. Members 18 – 59 years of age whose BP was <140/90 Members 60 – 85 years of age whose BP was <150/90 If a Diabetic, then 18-85 years of age whose BP was <140/90	76%
12	Prevention	CTC MSSP QIP (P4P)	Depression Screening	The percentage of patients 18 years and older seen two or more times within the last 24 months and who were screened one or more times for depression using a standardized screening tool (e.g. PHQ-2) during the measurement period.	75%
13	Prevention	BCBS CTC QIP (P4P)	Chlamydia Screening: Part A. Obtaining Sexual History	The percentage of women 16–24 years of age who were screened for sexual history during the measurement period.	59%
14	Prevention	BCBS CTC QIP (P4P)	Chlamydia Screening: Part B. Testing	The percentage of women 16–24 years of age who were identified as sexually active and who had at least one test for Chlamydia during the measurement period.	34%
15	Prevention	BCBS MSSP QIP (P4P) UHC Comm UHC MA	Breast Cancer Screening	The percentage of women 50-74 years of age who had a mammogram to screen for breast cancer during the measurement period or 18 months prior to the MP ¹ .	82%
16	Prevention	BCBS MSSP QIP (P4P) UHC Comm UHC MA	Colorectal Cancer Screening	The percentage of patients 50-75 years of age who had a colonoscopy during the measurement period or ten years prior to the MP ¹ .	72%
17	Prevention	UHC Comm	Cervical Cancer Screening	The percentage of patients 21- 64 years of age who received one or more Pap tests to screen for cervical cancer during the measurement period or 24 months prior to the MP ¹ .	85%
18	Prevention	BCBS UHC MA	Osteoporosis Management in Women who had a Fracture	The percentage of women 67 years of age and older who suffered a fracture during the MP period and who had either a bone mineral density (BMD) test or prescription for a drug to treat or prevent osteoporosis within six months of the fracture.	60%
19	Prevention	UHC MA	Rheumatoid Arthritis Management	The percentage of members 18 and older who were diagnosed with rheumatoid arthritis and who were dispensed at least one ambulatory prescription for a disease-modifying anti-rheumatic drug (DMARD).	78%
20	Care	MSSP	Medication Documentation	The percentage of members who had their medication list documented at EVERY visit in 2016.	N/A

¹ MP = Measurement Period

2016 Gaps in Care Adult Measures
QIP/CTC/BCBSRI/UHC/MSSP

	Condition	Payer	Measure	Description	Target
21	Prevention	MSSP	Influenza Immunization	The percentage of members 6 months or older in the medical record that contain documentation that the patient received influenza immunization between 10/2/2015 and 3/31/2016 or if there is any documentation that the immunization was given prior to 10/1/2015 but still in 2015.	N/A
22	Prevention	MSSP	Pneumonia Vaccination	The percentage of members 65 years and older in the medical record that indicate the patient has had the Pneumonia Vaccination (10 year look back)	N/A
23	Prevention	MSSP	High Blood Pressure Screening	The percentage of members 18 and older seen during the <i>MP</i> ¹ who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current BP reading as indicated If systolic ≥ 140 or diastolic > 90 patient should have a follow-up appointment in 1 month, if systolic is between 120-139 and diastolic is between 80-90, patient must be screened again in 1 year.	N/A
24	IVD	MSSP	Ischemic Vascular Disease	The percentage of members 18 years and older diagnosed with an Acute Myocardial Infarction (AMI), a coronary artery bypass graft (CABG) or had a percutaneous coronary interventions (PCI) in 2015 or had a diagnosis of Ischemic Vascular Disease (IVD) during 2016. If yes do they have a documented use of aspirin or other antithrombotic including Clopidogrel, Plavix, Prasugrel Effient or Ticlopidine .	N/A
25	Heart Failure	MSSP	Heart Failure Beta Blocker	The percentage of members 18 years and older with a diagnosis of heart failure AND a documented LVEF of <40 . If yes are they are on a beta blocker only including: Bisoprolol, Carvedilol or Metoprolol .	N/A
26	CAD	MSSP	Coronary Artery Disease ACE or ARB therapy for Patients with CAD and Diabetes and/or LVSD	The percentage of members 18 years and older with a diagnosis of CAD and either DM or LVSD (LVDS is an EF <40 . If yes is the patient on an ACE or ARB?	N/A
27	Mental Health	MSSP	Depression Remission at 12-months	The percentage of members 18 and older with a diagnosis of major depression who had an initial PHQ-9 score greater than 9 who show a remission at 11-13 months outward. Need to look at PHQ-9 score at visit in 2015 where it is greater than 9 and then look out 11-13 months into 2016 to see if PHQ-9 is now less than 5.	N/A

¹ MP = Measurement Period