

**2016 Quality Improvement Program (QIP) Adult Measures
BCBSRI/UHC**

	Condition	Measure	Description	Target
1	Diabetes	HbA1c Poor Control	The percentage of patients 18-75 years of age with diabetes whose most recent HbA1c level during the measurement period is >9.0% (poor control) or there is no HbA1c in the record.	≤20%
2	Diabetes	HbA1c Control	The percentage of patients 18-75 years of age with diabetes with the most recent HbA1c <8.0% (control) during the <i>MP</i> ¹ .	72%
3	Diabetes	Blood Pressure Control	The percentage of patients 18-75 with diabetes whose most recent blood pressure reading value is less than 140/90.	80%
4	Prevention	Tobacco Use	Patients age 18 years and older seen by a physician a minimum of once per measurement period who were queried about tobacco use a minimum of once per measurement period.	95%
5	Prevention	Tobacco Cessation Intervention	Patients age 18 years and older seen by a physician a minimum of once per measurement period who have been identified as smokers or tobacco users who received cessation intervention – (advice to quit, counseling, referral to counseling, and/or pharmacologic therapy during the measurement period).	85%
6	Prevention	Adult Body Mass Index (BMI)	The percentage of patients 18–74 years of age who had an outpatient visit and who had their “latest” body mass index (BMI) documented during the measurement period and whose BMI is either in the normal range OR is above or below the normal range and have a documented follow up plan within the measurement period.	87%
7	Prevention	Screening for Fall Risk	Percentage of patients aged 65 years and older who were screened for future fall risk at least once within the <i>MP</i> ¹ .	70%
8	Hypertension	Blood Pressure Control	The percentage of patients 18-85 years of age who had a diagnosis of hypertension (HTN) and their BP was adequately controlled during the measurement period. Members 18 – 59 years of age whose BP was <140/90 Members 60 – 85 years of age whose BP was <150/90 If a Diabetic, then 18-85 years of age whose BP was <140/90	76%
9	Prevention	Depression Screening	The percentage of patients 18 years and older seen two or more times within the last 24 months and who were screened one or more times for depression using a standardized screening tool (e.g. PHQ-2) during the measurement period.	75%
10	Prevention	Chlamydia Screening: Part A. Obtaining Sexual History	The percentage of women 16–24 years of age who were screened for sexual history during the measurement period.	59%
11	Prevention	Chlamydia Screening: Part B. Testing	The percentage of women 16–24 years of age who were identified as sexually active and who had at least one test for Chlamydia during the measurement period.	34%
12	Prevention	Breast Cancer Screening	The percentage of women 50-74 years of age who had a mammogram to screen for breast cancer during the <i>MP</i> ¹ or 18 months prior to the <i>MP</i> ¹ .	82%
13	Prevention	Colorectal Cancer Screening	The percentage of patients 50-75 years of age who had a colonoscopy during the <i>MP</i> ¹ or ten years prior to the <i>MP</i> ¹ .	72%

¹MP = Measurement Period